

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 11-07-2007		2. REPORT TYPE Final Report		3. DATES COVERED (From – To) 20 May 2006 - 09-Oct-08	
4. TITLE AND SUBTITLE Polydentate Halogen Bonding Donors for the Self-Assembly of New Materials			5a. CONTRACT NUMBER FA8655-06-1-3040		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Professor Giuseppe Resnati			5d. PROJECT NUMBER		
			5d. TASK NUMBER		
			5e. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Polytechnic of Milan 7, via Mancinelli Milan I-20052 Italy			8. PERFORMING ORGANIZATION REPORT NUMBER N/A		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) EOARD Unit 4515 BOX 14 APO AE 09421			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S) Grant 06-3040		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT This report results from a contract tasking Polytechnic of Milan as follows: When halogen bonding drives the self-assembly of new materials, the involved modules can be grouped into three main classes: XB donors, XB acceptors, XB self-complementary modules. The design of new XB donors was pursued in this project. Polydentate modules were pursued with the aim of having a better control on the self-assembled architectures they will form with XB acceptors. The high stereoselectivity of solid state reactions under the control of XB was further explored in other materials already prepared.					
15. SUBJECT TERMS Halogen bonding, self-assembly, Chemistry, EOARD					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UL	18, NUMBER OF PAGES 13	19a. NAME OF RESPONSIBLE PERSON BARRETT A. FLAKE
a. REPORT UNCLAS	b. ABSTRACT UNCLAS	c. THIS PAGE UNCLAS			19b. TELEPHONE NUMBER (Include area code) +44 (0)1895 616144



POLITECNICO DI MILANO

Dipartimento di Chimica, Materiali e Ingegneria Chimica "G. Natta"

Via Mancinelli 7 - 20131 Milano

Centralino: 02-23993000
Fax: 02-23993080
Segreteria amministrativa: 02-23993035
secretar@dept.chem.polimi.it
Codice Fiscale 80057930150
Partita IVA 04376620151

Prof. Dr. Giuseppe Resnati
ph.: +39-02-23993032
fax: +39-02-23993080
e-mail: giuseppe.resnati@polimi.it
URL: <http://nfmlab.chem.polimi.it/>

17 July 2007

POLYDENTATE HALOGEN BONDING DONORS FOR THE SELF-ASSEMBLY OF NEW MATERIALS

Grant/Cooperative Agreement No.: FA8655-06-1-3040


Performance Final Comprehensive Report 15 May 2006 – 14 May 2007

Part A: Tetra- and hexadentate halogen bonding donors.


In the first three months of activity of this cooperative agreement, we have studied the self-assembly of the tetratopic halogen bonding donor **5** with the structurally related tetratopic halogen bonding acceptor **1** (Scheme 1). This latter module was also self-assembled with the α,ω -diiodoperfluoroalkanes **2-4** presenting alkyl chains of different lengths.

The tetratopic acceptor **1** was prepared through a synthetic sequence similar to that optimized for the synthesis of **5** (reaction of pentaerythritol with 4-chloropyridine, Scheme 2). The rationale underlying the self-assembly of **1** and **5** was that both modules can adopt a tetrahedral conformation; thanks to a reciprocal self-induced fitting, this conformation was expected to be preferred over the other possible conformations in both modules and the construction of a diamondoid network was expected (Scheme 3). Indeed this was the case! The pairing of the complementary binding sites in **1** and **5** leads to the formation of a diamondoid net which presents a remarkable interpenetration in order to avoid the presence of huge voids. Specifically, two different sets of five diamondoid networks interpenetrate and form two separate 5-fold interpenetrated networks (Figure 1). Two such 5-fold interpenetrated networks further interpenetrate to give the overall crystal packing which thus presents a 10-fold interpenetration (Figure 2). This is a record for interpenetration in halogen bonded architectures.

Scheme 1



Scheme 2



Scheme 3





Figure 1


A 5+...-FOLD DIAMONDROID INTERPENETRATED NETWORK



N...I distance: 2.817 ?

N...I-C angle: 175.2°

5-fold network



View down the crystallographic a axis

Schematic view of the network





Figure 2



When the tetratopic module **1** was challenged with the ditopic modules **2-4**, the formed networks show a 1:2 ratio between the halogen bonding acceptor and donor modules and present complex interpenetrate assembly of different topologies. A rare diamondoid 8-fold network of class Ia¹ is formed when starting from 1,4-diiodotetrafluorobutane **2** (Scheme 4, Figure 3), while 2D square 4⁴ layers with 4-fold and 5-fold interpenetration are obtained when starting from 1,6-diiodoperfluorohexane **3** and from 1,8-diiodoperfluorooctane **4**, respectively (Scheme 4, Figure 4). .

Scheme 4




Figure 3





Figure 4



Besides studying bidentate and tetradentate halogen bonding donors (**2-4** and **5**, respectively), we also turned our attention to hexakis[4-(2,3,5,6-tetrafluoro-4-iodophenoxy)methyl]phenoxy]cyclotriphosphazene **6** which is the first hexadentate halogen bonding donor ever studied. The compound was prepared by reacting 4-hydroxy benzaldehyde with hexachlorocyclotriphosphazene in the presence of a base, reducing the carbonyls of the hexasubstituted product by using LiBH_4 at room temperature, and finally forming regioselectively the hexakis[4-(2,3,5,6-tetrafluoro-4-iodophenoxy)] derivative by reaction with iodopentafluorobenzene under basic conditions.

On evaporation of a solution containing 4,4'-dipyridylethylene, the phosphazene derivative **6** behaves as a nanopillar that self-assembles by halogen bonding with the telechelic partner and forms infinite supramolecular architectures (Scheme 5, Figure 5). Dipyridyl behaves in a strictly similar way (Figure 6). No interpenetration occurs as layered architectures are formed in both cases.

Scheme 5




Figure 5





Figure 6



Part B: Halogen bonding donors with eight or more binding sites

We have also investigated the synthesis of fluorinated poly(propylene imine) dendrimers (DAB-Am) dendrimers of the first, second, and third generation in order to prepare a new type of halogen bond donors.

The functionalisation of poly(propylene imine) dendrimers (DAB-Am) with perfluorinated groups was carried out by nucleophilic attack of the nitrogen at the periphery on different halopentafluorobenzenes and hexafluorobenzene (Scheme 5).



Scheme 5. Synthesis of dendrimeric tectons **3a-d**, **4a-d**, and **5b-d**.

The S_NAr attack of DAB-dendr-(NH₂)₂ⁿ⁺¹ (**1a-c**) on the fluorobenzenes **2a-d** afforded three different generations of DAB-dendr-(NHC₆F₄X)₂ⁿ⁺¹ [with n = 1 (**3**), n = 2 (**4**), n = 3 (**5**) and X = F (**a**), Cl (**b**), Br (**c**), I (**d**)] (Scheme 1). The first generation derivatives **3a-d** (n = 1,) have been obtained in 52 to 92% isolated yields after chromatographic purification (dichloromethane/methanol as eluent); the second (n = 2, **4a-d**) and the third generations (n = 3, **5b-d**), in 45 to 86% isolated yields. ¹⁹F NMR indicates that reactions invariably occur with high regioselectivity on the fluorine atom *para* to the heavy halogen of **2b-d**. Regioselectivity is largely unaffected by dendrimer size while it increases with the size of the heavy halogen (I > Br > Cl; the *ortho* by-product is 4% and 21% in **3d** and **3b**, respectively).


Clearly, the S_NAr reactions with **2c,d** transformed dendrimers **1**, which expose amine groups at their outer shells, into **3c,d-5c,d**, which expose C₆F₄-X groups, which are known to be strong XB-donor sites. The general ability of halotetrafluorophenyl substituted dendrimers to work as XB-based tectons was proven, in solution, by monitoring the high-field shift of the CF=CX signal in ¹⁹F NMR spectra on pyridine addition (see Table 1). This shift is diagnostic of the occurrence of XB and, as expected, the larger the amount of added pyridine, the greater the upfield shift is. Similar changes of chemical shifts and similar trends in the changes were also shown by second and third generation dendrimers **4d** and **5d**. The bromotetrafluorophenyl substituted dendrimers of first, second, and third generation **3c**, **4c**, and **5c**, respectively, also showed a similar behaviour. The only difference was that, for a given XB-donor/XB-acceptor ratio, shift changes were invariably smaller, consistent with the fact that bromopfluoroarenes are weaker XB-donors than iodopfluoroarenes.

Table 1. CF=CX signal shifts in ^{19}F NMR spectra of 3c,d-5c,d on pyridine addition.

Different amounts of excess pyridine were added to a 0.005 M solution of dendrimers **3c,d**, **4c,d**, and **5c,d** in CDCl_3 (CFCI_3 as internal standard). The upfield shift changes ($\Delta\delta$ (ppm) = $\delta(\text{pure dendrimer}) - \delta(\text{dendrimer in the presence of pyridine})$) for the CF=CX signals are reported in Table 2, while the CF-CF=CX shift changes, if any, were negligible. The amount of added pyridine changed with the generation (e.g. 50 equivalents for generation 1, 100 equivalents for generation 2, 200 equivalents for generation 3) so that the ratio between iodine atoms number and pyridine nitrogen atoms number (namely the ratio between XB-donor and XB-acceptor sites) was either 2/25 or 2/10 in all the experiments.

Dendrimer	I/N ratio	$\Delta\delta$ (ppm)
3c	2/25	0.03
4c	2/25	0.04
5c	2/25	0.05
3d	2/25	0.05
3d	2/10	0.02
4d	2/25	0.06
4d	2/10	0.03
5d	2/25	0.09
5d	2/10	0.04

In solution, ^{19}F NMR proves the involvement of iodine atoms of the DAB-*dendr*-($\text{NH-C}_6\text{F}_4\text{I}$) $_2^2$ **3d** in the formation of strong XB. In the solid, the four chains of **3d** look adjusted for an *exo* recognition process driven by XB. These observations prompted us to study the self-assembly of the DAB-*dendr*-($\text{NH-C}_6\text{F}_4\text{I}$) $_2^2$ **3d** with a telechelic nitrogen module, namely (*E*)-1,2-bis-(4-pyridil)-ethylene (**6**) (1:2 ratio, chloroform solution) (Scheme 6). Upon slow evaporation of the solvent, good quality crystals of **7** were deposited. ^1H NMR proved that the **3d/6** ratio in **7** is 1:2.

Scheme 6.

Few single X-ray structures of substituted DAB dendrimers are reported in the literature. For instance, it has been shown that chains of the DAB-*dendr*-(NH-Gly-t-BOC) $_4$ were backfolded due to the presence of hydrogen bonds between the CO and NH groups. After crystallisation in isopropyl ether, the structure of the DAB-*dendr*-($\text{NH-C}_6\text{F}_4\text{I}$) $_4$ **2a** has been determined through a single crystal X-ray diffraction (Figure 7). The analysis revealed that two opposite chains are fully extended in a *trans-trans* (*tt*) conformation despite the presence of two H...F hydrogen bonds. On the other hand, the other two chains are in a *gauche-gauche* conformation (*gg*), due to two N...H...F hydrogen bonds.




Figure 7: X-ray structure of DAB-dendr-(NHC₆F₄I)₄ **2a**.

In solution, ¹⁹F NMR proves the involvement of iodine atoms of the DAB-dendr-(NHC₆F₄I)₂ **3d** in the formation of strong XB. In the solid, the four chains of **3d** look adjusted for an *exo* recognition process driven by XB. These observations prompted us to study the self-assembly of the DAB-dendr-(NH-C₆F₄I)₂ **3d** with a telechelic nitrogen module, namely (*E*)-1,2-bis-(4-pyridil)-ethylene (**6**) (1:2 ratio, chloroform solution) (see Scheme 2, Supporting Information). Upon slow evaporation of the solvent, good quality crystals of **7** were deposited. ¹H NMR proved that the **3d**/**6** ratio in **7** is 1:2. Single crystal X-ray diffraction analysis showed that all of the DAB

Single crystal X-ray diffraction analysis showed that all of the DAB units function as tetratopic electron acceptors and the bipyridine derivatives act, in turn, as ditopic donors. XB is the key feature in the structure of **7** (Figure 8). The I1...N35[x,1+y,z] and I2...N29[1/2-x,-3/2+y,1/2-z] distances are 2.838 and 2.943 Å, around 0.8 times the sum of van der Waals radii for N and I.^[12] Consistent with the strong n→σ* character of the electron donation from nitrogen to iodine, the C-I...N angles that are almost linear (172.7 and 177.6°).




Figure 8. Single crystal X-ray structure of the supramolecular complex **7**; 2D square layers with a 4⁴ topology are formed.

Typical procedure for the reaction of DAB-dendr-(NH₂)₄ 1a with the perfluoroarene 2d.

A mixture of 100 mg (0.31 mmol) of DAB-dendr-(NH₂)₄, 0.33 mL (2.52 mmol) of pentafluoriodobenzene and 196 mg (1.42 mmol) of K₂CO₃ is stirred in 1 mL of refluxing CH₃CN for 24 h. Then the reaction is cooled to room temperature and the solid is filtered. After evaporation of the solvent, the crude material is chromatographed on silica gel with CH₂Cl₂ then CH₂Cl₂ / MeOH 93/7 as eluent.

DAB-dendr-(NH-C₆F₄I)₄ (C₄₀H₃₆N₆F₁₆I₄) 3d

White solid, mp = 100-103°C; 68% yield; IR ν_{\max} = 3424, 3165, 2953, 2823, 1640, 1524, 1488, 1149, 947, 796 cm⁻¹. ¹H NMR (250 MHz, CDCl₃): δ 5.03 (4H, l, NH), 3.47 (8H, m, NH-CH₂), 2.53 (8H, t, *J* = 6.4 Hz, CH₂-N), 2.42 (4H, m, CH₂-N), 1.75 (8H, qt, *J* = 6.4 Hz, CH₂-CH₂-NH), 1.45 (4H, m, N-CH₂-CH₂); ¹⁹F NMR (235 MHz, CDCl₃): δ -158.2 (8F, d, *J* = 18 Hz, CF-CN), -124.5 (8F, d, *J* = 18 Hz, CF-Cl); Ortho-substitution δ -168.3 (4F, m, CF-CF-CN), -156.6 (4F, m, CF-CN), -155.6 (4F, m, CF-CF-Cl), -115.3 (4F, m, CF-Cl); MS (ESI) *m/z* 1413 (M+H⁺).

DAB-dendr-(NH-C₆F₄Br)₄ (C₄₀H₃₆N₆F₁₆Br₄) 3c

White solid, mp = 85-87°C; 74% yield; IR ν_{\max} = 3423, 3220, 2954, 2822, 1640, 1518, 1494, 1154, 956, 821 cm⁻¹. ¹H NMR (250 MHz, CDCl₃): δ 4.89 (4H, l, NH), 3.45 (8H, m, NH-CH₂), 2.55 (8H, m, CH₂-N), 2.44 (4H, m, CH₂-N), 1.76 (8H, qt, *J* = 6.3 Hz, CH₂-CH₂-NH), 1.46 (4H, m, N-CH₂-CH₂); ¹⁹F NMR (235 MHz, CDCl₃): δ -158.2 (8F, d, *J* = 18 Hz, CF-CN), -137 (8F, d, *J* = 18 Hz, CF-CBr); Ortho-substitution δ -170.2 (4F, m, CF-CF-CN), -158.2 (4F, m, CF-CN), -157.4 (4F, m, CF-CF-CBr), -130.9 (4F, m, CF-CBr); MS (ESI) *m/z* 1225 (M+H⁺) most abundant peak of the isotope cluster of title compound.

DAB-dendr-(NH-C₆F₄Cl)₄ (C₄₀H₃₆N₆F₁₆Cl₄) 3b

White solid, mp = 73-75°C; 92% yield; IR ν_{\max} = 3434, 3212, 2951, 2812, 1645, 1520, 1496, 1149, 953, 871, 856 cm⁻¹; ¹H NMR (250 MHz, CDCl₃): δ 4.93 (4H, l, NH), 3.45 (8H, m, NH-CH₂), 2.53 (8H, t, *J* = 6.1 Hz, CH₂-N), 2.42 (4H, m, CH₂-N), 1.75 (8H, qt, *J* = 6.1 Hz, CH₂-CH₂-NH), 1.45 (4H, m, N-CH₂-CH₂); ¹⁹F NMR (235 MHz, CDCl₃): δ -159.9 (8F, d, *J* = 16.8 Hz, CF-CN), -144.7 (8F, d, *J* = 16.8 Hz, CF-CCl); Ortho-substitution δ -171.3 (4F, m, CF-CF-CN), -159.1 (4F, m, CF-CN), -158.8 (4F, m, CF-CF-CCl), -140.3 (4F, m, CF-CCl); MS (ESI) *m/z* 1047 (M+H⁺) most abundant peak of the isotope cluster of title compound.

DAB-dendr-(NH-C₆F₅)₄ (C₄₀H₃₆N₆F₂₀) 3a

White solid, mp = 73-75°C; 52% yield; IR ν_{\max} = 3440, 3210, 2963, 2823, 1662, 1520, 1483, 1018, 1000, 952, 800 cm⁻¹; ¹H NMR (250 MHz, CDCl₃): δ 4.60 (4H, l, NH), 3.38 (8H, m, NH-CH₂), 2.53 (8H, t, *J* = 6.5 Hz, CH₂-N), 2.43 (4H, m, CH₂-N), 1.74 (8H, qt, *J* = 6.5 Hz, CH₂-CH₂-NH), 1.45 (4H, m, N-CH₂-CH₂); ¹⁹F NMR (235 MHz, CDCl₃): δ -160.6 (8F, d, *J* = 22.0 Hz, CF-CN), -165.2 (8F, t, *J* = 21.4 Hz, CF), -173.0 (4F, t, *J* = 21.4 Hz, CF); MS (ESI) *m/z* 981 (M+H⁺).

Typical procedure for the reaction of DAB-dendr-(NH₂)₈ 1b with the perfluoroarene 2d.

A mixture of 100 mg (0.13 mmol) of DAB-dendr-(NH₂)₈, 0.30 mL (2.08 mmol) of pentafluoriodobenzene and 161 mg (1.17 mmol) of K₂CO₃ were stirred in 0.5 mL of refluxing THF during 48h. Then, the solution is filtered and the solvent is evaporated.

The crude material was chromatographed on silica gel eluent CH₂Cl₂ then CH₂Cl₂ / MeOH 8/2. (Syntheses were carried out in refluxing acetonitrile for **4a,b** and refluxing THF for **4c,d**).

DAB-dendr-(NH-C₆F₄I)₈ (C₈₈H₈₈N₁₄F₃₂I₈) 4d

Yellow oil; 65% yield; IR ν_{\max} = 3428, 2948, 2816, 1637, 1513, 1486, 1147, 947, 906, 800, 730 cm⁻¹; ¹H NMR (250 MHz, CDCl₃): δ 5.07 (8H, l, NH), 3.47 (16H, m, NH-CH₂), 2.53 (20H, m, N-CH₂-CH₂-CH₂-NH, N-CH₂-CH₂-CH₂-N), 2.41 (16H, m, N-CH₂-CH₂-CH₂-N, CH₂-CH₂-N), 1.75 (16H, qt, J = 5.9 Hz, CH₂-CH₂-NH), 1.61 (8H, m, N-CH₂-CH₂-CH₂-N), 1.34 (4H, m, N-CH₂-CH₂); ¹⁹F NMR (235 MHz, CDCl₃): δ -158.0 (2F, d, J = 19.9 Hz, CF-CN), -124.3 (2F, d, J = 19.9 Hz, CF-Cl) ; Ortho-substitution δ -168.4 (8F, m, CF-CF-CN), -156.7 (8F, m, CF-CN), -155.6 (8F, m, CF-CF-Cl), -115.3 (8F, m, CF-Cl) ; MS (ESI) m/z 2965-2966 (M+H⁺) isotope cluster.

DAB-dendr-(NH-C₆F₄Br)₈ (C₈₈H₈₈N₁₄F₃₂Br₈) 4c

Brown oil; 74% yield; IR ν_{\max} = 3423, 3220, 2950, 2817, 1640, 1492, 1151, 951, 823 cm⁻¹; ¹H NMR (250 MHz, CDCl₃): δ 4.90 (8H, l, NH), 3.46 (16H, m, NH-CH₂), 2.54 (20H, m, N-CH₂-CH₂-CH₂-NH, N-CH₂-CH₂-CH₂-N), 2.43 (16H, m, N-CH₂-CH₂-CH₂-N, CH₂-CH₂-N), 1.76 (16H, qt, J = 5.9 Hz, CH₂-CH₂-NH), 1.61 (8H, m, N-CH₂-CH₂-CH₂-N), 1.35 (4H, m, N-CH₂-CH₂); ¹⁹F NMR (235 MHz, CDCl₃): δ -159.1 (16F, d, J = 18.9 Hz, CF-CN), -137.0 (16F, d, J = 18.9 Hz, CF-CBr) ; Ortho-substitution δ -170.0 (8F, m, CF-CF-CN), -158.1 (8F, m, CF-CN), -157.4 (8F, m, CF-CF-CBr), -130.8 (8F, m, CF-CBr); MS (ESI) m/z 2589 (M +H⁺) most abundant peak of the isotope cluster of title compound.

DAB-dendr-(NH-C₆F₄Cl)₈ (C₈₈H₈₈N₁₄F₃₂Cl₈) 4b

Colourless oil; 86% yield; IR ν_{\max} = 3421, 3222, 2949, 2815, 1644, 1497, 1156, 957, 869, 739 cm⁻¹; ¹H NMR (250 MHz, CDCl₃): δ 4.94 (8H, l, NH), 3.45 (16H, m, NH-CH₂), 2.54 (20H, m, N-CH₂-CH₂-CH₂-NH, N-CH₂-CH₂-CH₂-N), 2.42 (16H, m, N-CH₂-CH₂-CH₂-N, CH₂-CH₂-N), 1.76 (16H, qt, J = 5.9 Hz, CH₂-CH₂-NH), 1.62 (8H, m, N-CH₂-CH₂-CH₂-N), 1.35 (4H, m, N-CH₂-CH₂); ¹⁹F NMR (235 MHz, CDCl₃): δ -159.9 (16F, m, CF-CN), -144.6 (16F, d, J = 17.7 Hz, CF-CCl) ; Ortho-substitution δ -171.2 (8F, m, CF-CF-CN), -159.2 (8F, m, CF-CN), -158.8 (8F, m, CF-CF-CCl), -140.2 (8F, m, CF-CCl) ; MS (ESI) m/z 2233 (M+H⁺) most abundant peak of the isotope cluster of title compound.

DAB-dendr-(NH-C₆F₅)₈ (C₈₈H₈₈N₁₄F₄₀) 4a

Colourless oil; 55% yield; IR ν_{\max} = 3423, 3242, 2949, 2814, 1640, 1517, 1482, 1018, 992, 962 cm⁻¹; ¹H NMR (250 MHz, CDCl₃): δ 4.62 (8H, l, NH), 3.38 (16H, m, NH-CH₂), 2.52 (20H, t, J = 6.3 Hz, N-CH₂-CH₂-CH₂-NH, N-CH₂-CH₂-CH₂-N), 2.39 (16H, m, N-CH₂-CH₂-CH₂-N, CH₂-CH₂-N), 1.74 (16H, qt, J = 6.3 Hz, CH₂-CH₂-NH), 1.58 (8H, m, N-CH₂-CH₂-CH₂-N), 1.33 (4H, m, N-CH₂-CH₂); ¹⁹F NMR (235 MHz, CDCl₃): δ -160.6 (16F, d, J = 22.0 Hz, CF-CN), -165.2 (16F, t, J = 22.0 Hz, CF), -173.2 (5F, d, J = 18.4 Hz, CF), -173.3 (3F, d, J = 16.9 Hz, CF) ; MS (ESI) m/z 2101-2102 (M+H⁺) isotope cluster.

Typical procedure for the reaction of DAB-dendr-(NH₂)₁₆ 1c with the perfluoroarene 2d.

A mixture of 100 mg (0.06 mmol) of DAB-dendr-(NH₂)₁₆, 0.25 mL (1.90 mmol) of pentafluoriodobenzene and 141 mg (1.02 mmol) of K₂CO₃ were stirred in 0.5 mL of

refluxing THF during 48h. Then, the solution is filtered and the solvent is evaporated. The crude material was chromatographed on silica gel eluent CH₂Cl₂ then CH₂Cl₂ / MeOH 7/3.

DAB-dendr-(NH-C₆F₄I)₁₆ (C₁₈₄H₁₉₂N₃₀F₆₄I₁₆) 5d

Yellow oil; 50% yield; ¹H NMR (250 MHz, CDCl₃): δ 5.10 (16H, I, NH), 3.46 (32H, m, NH-CH₂), 2.52 (56H, m, N-CH₂-CH₂-CH₂-NH, N-CH₂-CH₂-CH₂-N), 2.40 (28H, m, N-CH₂-CH₂-CH₂-N, CH₂-CH₂-N), 1.75 (32H, qt, *J* = 5.9 Hz, CH₂-CH₂-NH), 1.58 (24H, m, N-CH₂-CH₂-CH₂-N), 1.25 (4H, m, N-CH₂-CH₂); ¹⁹F NMR (235 MHz, CDCl₃): δ -157.9 (2F, d, *J* = 19.0 Hz, CF-CN), -124.3 (2F, d, *J* = 19.0 Hz, CF-Cl); Ortho-substitution δ -168.4 (8F, m, CF-CF-CN), -156.7 (8F, m, CF-CN), -155.6 (8F, m, CF-CF-Cl), -115.3 (8F, m, CF-Cl); MS (ESI) *m/z* 2024 (M+3H)³⁺ isotope cluster not resolved.

DAB-dendr-(NH-C₆F₄Br)₁₆ (C₁₈₄H₁₉₂N₃₀F₆₄Br₁₆) 5c

Brown oil; 45% yield; ¹H NMR (250 MHz, CDCl₃): δ 5.03 (16H, I, NH), 3.44 (32H, m, NH-CH₂), 2.54 (56H, m, N-CH₂-CH₂-CH₂-NH, N-CH₂-CH₂-CH₂-N), 2.44 (28H, m, N-CH₂-CH₂-CH₂-N, CH₂-CH₂-N), 1.93-1.42 (56H, m, CH₂-CH₂-N), 1.30 (4H, m, N-CH₂-CH₂); ¹⁹F NMR (235 MHz, CDCl₃): δ -159.1 (32F, d, *J* = 18.3 Hz, CF-CN), -137.1 (32F, d, *J* = 18.3 Hz, CF-CBr); Ortho-substitution δ -170.2 (16F, m, CF-CF-CN), -158.2 (16F, m, CF-CN), -157.4 (16F, m, CF-CF-CBr), -130.9 (16F, m, CF-CBr); MS (ESI) *m/z* 2660 (M+2H)²⁺ isotope cluster not resolved.

DAB-dendr-(NH-C₆F₄Cl)₁₆ (C₁₈₄H₁₉₂N₃₀F₆₄Cl₁₆) 5b

Colourless oil; 51% yield; ¹H NMR (250 MHz, CDCl₃): δ 4.99 (16H, I, NH), 3.44 (32H, m, NH-CH₂), 2.52 (56H, m, N-CH₂-CH₂-CH₂-NH, N-CH₂-CH₂-CH₂-N), 2.37 (28H, m, N-CH₂-CH₂-CH₂-N, CH₂-CH₂-N), 1.75 (32H, m, CH₂-CH₂-NH), 1.59 (24H, m, N-CH₂-CH₂-CH₂-N), 1.32 (4H, m, N-CH₂-CH₂); ¹⁹F NMR (235 MHz, CDCl₃): δ -159.9 (32F, d, *J* = 18.3 Hz, CF-CN), -144.7 (32F, m, CF-CCl); Ortho-substitution δ -171.3 (16F, m, CF-CF-CN), -159.2 (16F, m, CF-CN), -158.8 (16F, m, CF-CF-CCl), -140.3 (16F, m, CF-CCl); MS (ESI) *m/z* 2304 (M+2H)²⁺ isotope cluster not resolved.

Characterization of supramolecular complex 7:

White solid, mp = 162-165°C; IR *v*_{max} (cm⁻¹) = 3414, 3186, 2950, 2873, 2816, 1639, 1598, 1488, 1290, 1140, 944, 782 cm⁻¹.

The principal investigator

Prof. Giuseppe Resnati

